

Division of Environmental Health and Communicable Disease Prevention		
Section: 4.0 Diseases and Conditions	Updated 3/05	

Page 1 of 12

# Psittacosis Table of Contents

Psittacosis
Fact Sheet
Disease Case Report (CD-1)
Record of Investigation of Communicable Disease (CD-2)

Subsection: Psittacosis



Division of Environmental Health and Co	ommunicable Disease Prevention
---	--------------------------------

Section: 4.0 Diseases and Conditions	Updated 3/05
Subsection: Psittacosis	Page 2 of 12

#### **Psittacosis**

# Overview (1,2)

For a complete description of psittacosis, refer to the following texts:

- Control of Communicable Diseases Manual (CCDM).
- Red Book, Report of the Committee on Infectious Diseases.

## **Case Definition** (3)

#### Clinical description

An illness characterized by fever, chills, headache, photophobia, cough, and myalgia

### Laboratory criteria for diagnosis

- Isolation of *Chlamydophila psittaci* (formerly *Chlamydia psittaci*) from respiratory secretions, or
- Fourfold or greater increase in antibody against *C. psittaci* by complement fixation or microimmunofluorescence (MIF) to a reciprocal titer of ≥32 between paired acute- and convalescent-phase serum specimens, or
- Presence of immunoglobulin M antibody against *C. psittaci* by MIF to a reciprocal titer of ≥16

#### Case classification

Confirmed: a clinically compatible case that is laboratory confirmed *Probable*: a clinically compatible case that is epidemiologically linked to a confirmed case or that has supportive serology (e.g., C. psittaci titer of  $\geq 32$  in one or more serum specimens obtained after the onset of symptoms)

#### Comment:

The serologic findings by CF may also occur as a result of infection with *Chlamydia pneumoniae* or *Chlamydia trachomatis*. The MIF is more sensitive and specific than CF for infection with *C. psittaci*, but availability of this newer test may be more limited.

# **Information Needed for Investigation**

- Verify the diagnosis. Determine what laboratory tests were conducted and the results.
- **Establish the extent of illness**. Determine if household or other close contacts are, or have been, ill by contacting the health care provider, patient or family member.
- Determine if a potential reservoir of infection exists in your community that may produce more cases.

Missouri Department of Health and Senior Services Communicable Disease Investigation Reference Manual



vision of Environmental Health and Communicable Disease Prevention			
Section: 4.0 Diseases and Conditions	Updated 3/05		
Subsection: Psittacosis	Page 3 of 12		

- Contact the Regional Communicable Disease Coordinator if an outbreak is suspected, or if cases are in high-risk settings such as child care, a medical facility, or long term care.
- Contact the Bureau of Child Care (573-751-2450) if cases are associated with a childcare facility.
- Contact the Section for Long-term Care Regulations (573-526-0721) if cases are associated with a long-term care facility.
- Contact the Bureau of Health Facility Regulation (573-751-6303) if cases are associated with a hospital or hospital-based long-term care facility.

#### **Notification And Control Measures**

Determine the source of infection.

- Psittacosis is rarely transmitted person-to-person. Most human cases may be traced back to contact with infected birds. Infected birds may exhibit overt signs of illness or be asymptomatic. Parakeets, parrots, and cockatiels have been the primary reservoir of infection but canaries, finches, pigeons, doves, chickens, turkeys, and many other species of birds may become infected.
- Ask about the patient's occupation, name, and location of employer. Interview the patient about job duties, if it appears the individual may have contracted the disease at work. Ask about patient contact with pet birds, live poultry, and other activities that would have exposed the individual to airborne dust contaminated with bird droppings or respiratory secretions.
- Identify symptomatic household members, associates, or co-workers and urge them to contact their physician for a medical evaluation.
- Although there is no epidemiologic evidence of increased risk to the very young, the elderly, or the immunocompromised, more rigorous testing should be considered for birds in contact with these individuals due to the potentially severe nature of this disease in humans.
- Individuals who have suspect, sick, or dying birds or poultry should consult a veterinarian. The Regional Communicable Disease Coordinator will contact the State Public Health Veterinarian who will contact the Missouri Department of Agriculture, as needed.
- It is imperative that individuals in contact with infected birds or contaminated materials take precautions to prevent infection, as described in the "*Chlamydophila* Compendium." (4) Precautions include the use of protective clothing, gloves, a disposable surgical cap, and an appropriately fitted respirator with N95 or higher rating. Information pertaining to respirator use can be obtained from the National Institute of Occupational Safety and Health: Safety and Health Topics, Respirators (http://www.cdc.gov/niosh/npptl/topics/respirators/ (March 2005)



I	Division of Environmental Health and Communicable Disease Prevention
ı	

Section: 4.0 Diseases and Conditions	Updated 3/05		
Subsection: Psittacosis	Page 4 of 12		

#### **Control Measures**

See the Control of Communicable Diseases Manual, Psittacosis, "Methods of control."

See the Red Book, Chlamydia (Chlamydophila) psittaci, "Control Measures."

See the <u>Compendium of Measures To Control Chlamydophila psittaci</u> (formerly <u>Chlamydia psittaci</u>) Infection Among Humans (Psittacosis) and Pet Birds, 2004. (4)

If the epidemiological data indicate a specific source of infection and the potential for ongoing transmission to humans, animal quarantine measures may be established. Animal quarantines or modified quarantines may be established by the local health authority, the Missouri Department of Health and Senior Services, and/or the Missouri Department of Agriculture.

### **Laboratory Procedures**

## **Specimens:**

Diagnosis of human cases is serological, based on a rising antibody titer between acute and convalescent sera specimens collected two to three weeks apart. Diagnostic testing is available through private laboratories. Cross reactions may occur with *C. pneumoniae* or *C. trachomatis* infections.

Confirmation by isolation of the organism from respiratory secretions is rarely attempted. Attempts to culture the organism in laboratories without specialized equipment has lead to outbreaks among laboratory personnel.

The Missouri State Public Health Laboratory (SPHL) no longer performs laboratory testing for psittacosis. Acute and convalescent sera specimens may be sent to CDC in special circumstances or for epidemiological purposes. Specimens should not be collected without prior authorization by the Regional Communicable Disease Coordinator. They will be sent through the SPHL to CDC. Please advise the submitter that serum specimens are accepted for epidemiological purposes but not for diagnosis and frequently several months may elapse before results are received. Additional information on laboratory procedures can be obtained from the Regional Communicable Disease Coordinator or from staff at the SPHL. The SPHL telephone number is 573-751-0633. The web site is <a href="http://www.dhss.mo.gov/Lab/index.html">http://www.dhss.mo.gov/Lab/index.html</a> (March 2005)

The University of Missouri's Veterinary Medical Diagnostic Laboratory (VMDL) in Columbia offers testing for *C. psittaci* infection/harborage of birds. Interested individuals and their veterinarians may contact the VMDL for a schedule of fees and specimen shipping requirements. The telephone number of the VMDL is 1-800-862-8635 or 573-882-6811. The web site is <a href="http://www.cvm.missouri.edu/vmdl">http://www.cvm.missouri.edu/vmdl</a>. (March 2005)

SOURI DEPARTMENT	Division of Environmental Health and Communica	able Disease Prevention
dhss &	Section: 4.0 Diseases and Conditions	Updated 3/05
TH AND SENIOR SERV	Subsection: Psittacosis	Page 5 of 12

## **Reporting Requirements**

Psittacosis is a Category II disease and shall be reported to the local health authority or to the Missouri Department of Health and Senior Services within three days of first knowledge or suspicion by telephone, facsimile or other rapid communication.

- 1. For all cases complete a "Disease Case Report" (CD-1).
- 2. For confirmed and probable cases complete a "Record of Investigation of Communicable Disease" (CD-2).
- 3. Entry of the completed CD-1 into MOHSIS negates the need for the paper CD-1 to be forwarded to the Regional Health Office.
- 4. Send the completed secondary investigation form to the Regional Health Office.
- 5. All outbreaks or "suspected" outbreaks must be reported as soon as possible (by phone, fax or email) to the Regional Communicable Disease Coordinator. This can be accomplished by completing the Missouri Outbreak Surveillance Report (CD-51).
- 6. Within 90 days of the conclusion of an outbreak, submit the final outbreak report to the Regional Communicable Disease Coordinator.

#### References

- Control of Communicable Diseases Manual. "Psittacosis (*Chlamydia psittaci* infection, Ornithosis, Parrot fever, Avian chlamydiosis)." Heymann, David L., ed 18<sup>th</sup> ed. Washington, DC: American Public Health Association, 2004: 432-434.
- 2. American Academy of Pediatrics. "*Chlamydia (Chlamydophila) psittaci* (Psittacosis, Ornithosis)." In: Pickering LK., ed. *Red Book: 2003 Report of the Committee on Infectious Diseases.* 26<sup>th</sup> ed. Elk Grove Village, IL. American Academy of Pediatrics; 2003: 237-238.
- 3. Centers for Disease Control and Prevention. *Case Definitions for Infectious Conditions Under Public Health Surveillance*. MMWR 1997: 46 (No. RR-10). "Psittacosis (*Chlamydia psittaci*) (Ornithosis)," 1996, <a href="http://www.cdc.gov/epo/dphsi/casedef/psittacosiscurrent.htm">http://www.cdc.gov/epo/dphsi/casedef/psittacosiscurrent.htm</a>. (March 2005)
- 4. National Association of State Public Health Veterinarians, <u>Compendium of Measures To Control Chlamydophila psittaci</u> (formerly <u>Chlamydia psittaci</u>) Infection Among Humans (<u>Psittacosis</u>) and <u>Pet Birds</u>, <u>2004</u>. <a href="http://www.avma.org/pubhlth/psittacosis.asp">http://www.avma.org/pubhlth/psittacosis.asp</a>. (March 2005)

# **Other Sources of Information**

- 1. Mandell, Douglas and Bennett's *Principles and Practice of Infectious Diseases*. "*Chlamydia psittaci* (Psittacosis)." G. Mandell, J. Bennett, R. Dolin, eds. 6<sup>th</sup> ed. Vol. 2, 2005: 2251, 2256-2258.
- 2. Bacterial Infections of Humans Epidemiology and Control. "Chlamydial Infections." Schachter, Julius, and Alexander, E. Russell. 3<sup>rd</sup> ed. Eds. Alfred S. Evans and Philip S. Brachman. New York: Plenum, 1998: 197-222.

Missouri Department of Health and Senior Services Communicable Disease Investigation Reference Manual



Division of Environmental Health and Communication	able Disease Prevention
Section: 4.0 Diseases and Conditions	Updated 3/05
Subsection: Psittacosis	Page 6 of 12

- 3. The Merck Veterinary Manual. 8<sup>th</sup> Ed. Ed. Susan E. Aiello. Whitehouse Station, NJ: Merck & Co., Inc., 1998. <a href="http://www.merckvetmanual.com/mvm/index.jsp">http://www.merckvetmanual.com/mvm/index.jsp</a> (March 2005) (search "psittacosis" or "psittaci").
- 4. Zoonoses and Communicable Diseases Common to Man and Animals. "Avian Chlamydiosis." 2<sup>nd</sup> ed. Eds. Pedro Acha and Boris Szyfres. Pan American Health Organization, Washington, DC, Scientific Publication No. 503, 1994: 250-255.

# **Web Resources and Information**

- 1. Psittacosis Fact Sheet, Maryland Department of Health (several languages available) http://www.edcp.org/factsheets/psittfact.html. (March 2005)
- 2. Psittacosis, Farhad Arjomand, & Lessnau, K.D., eMedicine Journal, May 3 2002, V 3, N 5 <a href="http://www.emedicine.com/med/topic1951.htm">http://www.emedicine.com/med/topic1951.htm</a> (March 2005)

#### **Psittacosis**

**FACT SHEET** 

#### What is psittacosis?

Psittacosis is a bacterial disease also referred to as parrot fever or ornithosis. The agent that causes this disease is *Chlamydophila psittaci*. The symptoms of the disease are variable and may include fever, rash, muscle aches, chills, and a dry cough. It may produce pneumonia, which may not be apparent except on x-ray. The disease is transmitted from birds to people. Parakeets, parrots, and cockatiels are among the main carriers of infection. All birds are susceptible to the disease and outbreaks have occurred on poultry farms, in pet shops, and in processing plants.

## Who gets psittacosis?

Anyone can get psittacosis if they are exposed to infected birds. Occupations at highest risk in the United States are bird breeders, poultry processing workers, farmers who raise poultry, veterinarians, pet shop owners and their employees. Sporadic cases also occur in individual households that have pet birds.

## How do humans get psittacosis?

The infection is acquired by inhaling the agent in dusts from dried respiratory secretions and droppings of infected birds.

## How would I know if my birds have psittacosis?

Birds may be harboring the bacteria without any apparent ill effects. When birds develop the disease they become quiet, withdrawn, lose weight, and exhibit a pronounced change in their feces or droppings. If you believe your pet birds or poultry may be infected, you should contact your veterinarian who can arrange for the appropriate laboratory testing.

# What are the symptoms of psittacosis in people?

Fever, muscle aches, chills, headache, and a dry cough. A rash may also be present.

# How long is the incubation period for psittacosis?

The incubation period is one to four weeks.

# How is psittacosis diagnosed?

Since the disease is uncommon in the United States, the diagnosis usually requires laboratory tests. Several different methods are available from commercial laboratories to aid the clinician in diagnosing the disease. Exposure history is very important for the diagnosis.

# What is the treatment for psittacosis?

Antibiotics such as tetracycline or erythromycin for 10 to 14 days are usually sufficient.

#### Can human to human transmission of psittacosis occur?

If one person has pneumonia caused by this agent and he/she coughs, the infection can theoretically spread to other people. However, this is an unlikely form of transmission because the cough is usually non-productive and the ill individual normally does not expel the bacteria from the lungs.

## What possible complications may arise from an untreated infection?

Encephalitis, meningitis, endocarditis, and neurological complications may occur. Severe pneumonia or death may occur in the elderly or immunocompromised.

#### How can psittacosis be prevented?

The most important measures to control the disease are already in place in the United States. For the most part, commercial importation of psittacine (parrot and parrot-like) birds ended in 1993 with the implementation of the Wild Bird Conservation Act. Persons who import birds must comply with federal laws that include regulatory requirements such as mandatory quarantines and prophylactic antibiotic treatment. Ongoing programs conducted by the state departments of agriculture provide for prevention of the spread of the disease in domestic poultry.

Consumers should purchase all birds only from legitimate retailers. Most pet shops, hatcheries, and retailers maintain records that aid in the traceback of infection should a case arise in one or more birds. This helps ensure that birds are treated, as are any persons who have handled or purchased the birds.

Individuals who may have contact with infected birds or contaminated materials should take precautions to prevent infection, such as wearing protective clothing, gloves, a disposable surgical cap, and an appropriately fitted respirator with N95 or higher rating. Information pertaining to respirator use can be obtained from the National Institute of Occupational Safety and Health: Safety and Health Topics, Respirators (<a href="http://www.cdc.gov/niosh/topics/respirators">http://www.cdc.gov/niosh/topics/respirators</a>). (March 2005)

Missouri Department of Health and Senior Services Section for Communicable Disease Prevention Phone: (866) 628-9891 or (573) 751-6113

#### MISSOURI DEPARTMENT OF HEALTH AND SENIOR SERVICES REPORT TO LOCAL PUBLIC HEALTH AGENCY DISEASE CASE REPORT 1 DATE OF REPORT 2 DATE RECEIVED BY LOCAL HEALTH AGENCY 3 NAME (LAST, FIRST, M.I.) 4 GENDER 5 DATE OF BIRTH 6 AGE 7 HISPANIC ☐ YES ☐ MALE ☐ FEMALE ☐ UNKNOWN 8 RACE (CHECK ALL THAT APPLY) 9 PATIENT'S COUNTRY OF ORIGIN 10 DATE ARRIVED IN USA ☐ BLACK ☐ ASIAN ☐ PACIFIC ISLANDER AMERICAN INDIAN □ WHITE ☐ UNKNOWN 11 ADDRESS (STREET OR RFD, CITY, STATE, ZIP CODE) 12 COUNTY OF RESIDENCE 13 TELEPHONE NUMBER 14 PREGNANT ☐ YES (IF YES NUMBER OF WEEKS 15 PARENT OR GUARDIAN 16 RECENT TRAVEL OUTSIDE OF MISSOURI OR USA 17 DATE OF RETURN ☐ YES ☐ NO ☐ UNKNOWN □ NO IF YES, WHERE 19 SCHOOL/DAY CARE/WORKPLACE 18 OCCUPATION ADDRESS (STREET OR RFD, CITY, STATE, ZIP CODE) 20 WORK TELEPHONE NUMBER 24 PATIENT RESIDE IN NURSING HOME 25 PATIENT DIED OF THIS ILLNESS 26 CHECK BELOW IF PATIENT OR 23 WAS PATIENT HOSPITALIZED PATIENT HHLD MEMBER MEMBER OF PATIENT'S ☐ YES ☐ NO ☐ UNKNOWN ☐ YES ☐ NO ☐ UNKNOWN ☐ YES ☐ NO ☐ UNKNOWN HOUSEHOLD (HHLD): NO UNK YES NO UNK 27 NAME OF HOSPITAL/NURSING HOME IS A FOOD HANDLER 28 HOSPITAL/NURSING HOME ADDRESS (STREET OR RFD, CITY, STATE, ZIP CODE) ATTENDS OR WORKS AT A CHILD OR ADULT DAY CARE CENTER 29 REPORTER NAME 30 TELEPHONE NUMBER IS A HEALTH CARE WORKER 31 REPORTER ADDRESS (STREET OR RFD, CITY, STATE, ZIP CODE) 32 TYPE OF REPORTER/SUBMITTER ☐ PHYSICIAN ☐ OUTPATIENT CLINIC ☐ PUBLIC HEALTH CLINIC ☐ HOSPITAL ☐ LABORATORY ☐ SCHOOL ☐ OTHER. 33 ATTENDING PHYSICIAN/CLINIC NAME ADDRESS (STREET OR RFD, CITY, STATE, ZIP CODE) **34** TELEPHONE NUMBER 35 DISEASE NAME(S) 36 ONSET DATE(S) 37 DIAGNOSIS DATE(S) 38 DISEASE STAGE/ 39 PREVIOUS DISEASE/STAGE 40 PREVIOUS DISEASE DATE(S) RISK FACTOR TEST DATE QUALITATIVE / COLLECTION DATE REFERENCE LABORATORY NAME/ADDRESS TYPE OF TEST SPECIMEN TYPE QUANTITATIVE RESULTS (MO/DAY/YR) RANGE (INCLUDE STREET OR RFD, CITY, STATE, ZIP CODE) - DIAGNOSTICS TREATED REASON NOT TREATMENT DATE TREATMENT DURATION PREVIOUS LOCATION **TREATMENTS** TYPE OF TREATMENT DRUG DOSAGE PREVIOUS TREATMENT Y/N/UNK) (MO/DAY/YR) (IN DAYS) (LIST CITY, STATE) 42 SYMPTOM ONSET DATE SYMPTOM DURATION SYMPTOM (IF APPLICABLE) SYMPTOM SITE (IF APPLICABLE) (MO/DAY/YR) (IN DAYS) SYMPTOMS 44 COMMENTS

#### **NOTES FOR ALL RELEVANT SECTIONS:**

- Stages, risk factors, diagnostics, treatments, and symptoms shown below are examples. To see a more complete listing, please go to
   http://www.dhss.state.mo.us/Diseases/DDwelcome.htm.

   You may also contact the Office of Surveillance at 1-800-392-0272 for additional information or to report a case.
- All dates should be in Mo/Day/Year (01/01/2001) format.
- All complete addresses should include city, state and zip code.
- · Required fields referenced below are italicized and bold, however fill form as complete as possible.
- (1) Date of Report -- date sent by submitter of document.
- (2) Date received will be filled in by receiving agency.
- (3-8) CASE DEMOGRAPHICS/IDENTIFIERS: Last name, First Name, Gender, Date of Birth, Hispanic, Race please check all that apply
- (23) Was patient hospitalized due to this illness?
- (32) Type of reporter/submitter (doctor, nursing home, hospital, laboratory) (33-34) Attending physician or clinic (full physician name and degree, address, phone)

Healthcare worker Converter/2 yrs  $\geq$  10 Converter/2 yrs  $\geq$  15

#### DISEASE: (35) Disease name or name(s), (36) Onset date(s), (37) Diagnosis Date(s)

#### (38) Disease Stage or Risk Factor

**Syphilis** Gonorrhea or Chlamydia **TB** Infection Primary (chancre present) Asymptomatic Contact to TB case Secondary (skin lesions, rash) Uncomplicated urogenital (urethritis, Immunocompromised Early Latent (asymptomatic < 1 year) cervicitis) Abnormal CXR Late Latent (over 1 year duration) Salpingitis (PID) Foreigner/Immigrant Neurosyphilis Ophthalmia/conjunctivitis IV Drug/Alcohol Abuse Cardiovascular Other (arthritis, skin lesions, etc) Resident, correctional Congenital Employee, correctional Other Over 70 Homeless Diabetes

(39) Previous Disease/Stage (if applicable) (40) Previous Disease Dates (if applicable)

#### (41) Diagnostics (Please Attach Lab Slip)

**Test Type** 

**Hepatitis** TB Other Igm Anti-HBc Not Done Elisa Anti-HBs Western Blot Mantoux Anti-HBc Total Multiple puncture device Culture Igm Anti-HAV ALT X-Ray HBsAa Smear AST Hep C Culture

Specimen Type (blood, urine, CSF, smear, swab), Collection Date (Mo/Day/Yr), Qualitative (negative, positive, reactive), Quantitative Results (1:1, 2.0 mm reading,) Reference Range (1:1neg, 1:64 equivocal, 1:128 positive, > 2 positive), Laboratory (name, address)

#### (42) TREATMENT

Reason not treated Drug
False positive TB
Previous treated Isoniazid
Age Ethambutol
Pyrazinamide
Rifampin

#### (43) SYMPTOMS:

**Symptom** (jaundice, fever, dark urine, headache) **Symptom Site** (head, liver, lungs, skin), **Symptom Onset Date** (Mo/Day/Yr) and **Symptom Duration** (in days)

(44) Comments: Attach additional sheets if more comments needed.

MO 580-0779 (9-01)

## MISSOURI DEPARIMENT OF HEALTH AND SENIOR SERVICES

# RECORD OF INVESTIGATION OF COMMUNICABLE DISEASE\*

Patient's Name				en like ilike negative e			ODING ONLY
						County	City
Address		City	State	Zip C	Code	Twnshp.	Disease
	I DF DW D	N 🗆 Other	County of Resi	dence		Hospital	Sour
arent's Name If Not	Adult		Phone	)			.
Hospitalized Ho	ospital Name			Date of Onset		Physician	5 8
Physician's Name						Phone Number	
Address	***************************************		-		Date		
Previous Address (if	significant)				Date	Moved	
Place Employed or So	chool Attended		7	Occupation		···	
Date Reported	How did you first 1	earn of this ca	ase?	1		Date	
Chief Clinical Sympto	oms with Dates:						
mier Cimical Sympto	oms with Dates:	and the same of the	J. Buther and Ava				7
						*	
1000		roduka da					
Treatment (type, amo	unt, dates):				•		
Type of Specimen	Date Collected	DIAGNOST	IC LABORATORY	TESTS ON PAT	IENT		
Type of Specimen	Date Collected		Result			Name of Labora	tory
				uso-			
			gerbin				
Are there other assoc	ciated cases?		_ If yes, how many	, and how associ	ated?		
							-
Household Sanitation	Good	Milk Supply					
	Poor	Water Suppl	у				
		(	Continued on res	verse side)			

to district the desired that the second state of the second						
						3
					2	"
			(Household and O	1	Т	
Name and Address	Age Sex	Relation to Patient	Similar Illness? Onset Date	Laboratory Specimen	Date Collected	Result
	č.					
United Till						
National National						
rative and Follow-up Notes						F 2
						2
HANDS SHOWN						
bable Source						
Recovered Died	Date of Deat	h	Cause of I	Peath		
vestigated by						